



Back to results | 1 of 1

View at Publisher | Export | Download | Add to List | More...

Malaysian Journal of Analytical Sciences

Volume 20, Issue 2, 2016, Pages 238-246

Two isolated crystalline solids from the stem of entada spiralis ridl. (Akar sintok)

(Article)

[Pemencilan dua jenis pepejal kristal dari batang pokok entada spiralis (Akar sintok)]

Harun, A.^a , So'ad, S.Z.M.^b , Hassan, N.M.^b

^a Faculty of Applied Sciences, Universiti Teknologi MARA Pahang, Bandar Tun Razak Jengka, Pahang, Malaysia

^b Kulliyah of Pharmacy, International Islamic University Malaysia, Bandar Indera Mahkota, Kuantan, Pahang, Malaysia

View references (23)

Abstract

Entada spiralis Ridl. (Leguminosae) is a woody climber which locally known as 'akar sintok'. The isolation and structure determination of two crystalline solids from active methanol fraction of E. spiralis stem bark had been examined. The structure identification of both solids was based on spectroscopic data (¹H and ¹³C NMR, HMQC, HMBC, DEPT135 and MS) along with comparison with literature data. The results revealed that these two crystalline solid sugars isolated from the most active fraction were known as penta-2-acetoxy-β-D-digitoxopyranosyl-(1→2)-fructofuranosyl-(6→4)-β-D-glucopyranosyl-(1→4)-glucopyranosyl-(1→2)-β-D fructofuranosyl-(6→1)-β-D-glucopyranosyl-(4→1)-acetylglucosamine (1) and β,D-glucopyranosyl-(1→2)-β,D-glucopyranosyl-(1→3)-β,D-xylopyranosyl-(1→4)-α,L-rhamnopyranosyl-(1→3)-β,D-glucopyranosyl-(1→3)-β,D-glucopyranoside (2) Thus, this finding can be utilized as a scientific baseline information for further skin disease studies, since both constituents were isolated from moderate antidermatophytic active fraction. © 2016, Malaysian Society of Analytical Sciences. All rights reserved.

Author keywords

Crystalline saponin; Entada spiralis; Glycosides; Isolation

ISSN: 13942506 Source Type: Journal Original language: English

Document Type: Article

Publisher: Malaysian Society of Analytical Sciences

View in search results format

References (23)

☐ All ☐ Export | ☐ Print | ☐ E-mail | ☐ Save to PDF | ☐ Create bibliography

☐ Mbosso, E.J.T., Ngouela, S., Nguedia, J.C.A., Beng, V.P., Rohmer, M., Tsamo, E.
1 **In vitro antimicrobial activity of extracts and compounds of some selected medicinal plants from Cameroon**
(2010) *Journal of Ethnopharmacology*, 128 (2), pp. 476-481. [Cited 35 times](#).
doi: 10.1016/j.jep.2010.01.017
[View at Publisher](#)

☐ Prasad, C.S., Shukla, R., Kumar, A., Dubey, N.K.
2 **In vitro and in vivo antifungal activity of essential oils of Cymbopogon martini and Chenopodium ambrosioides and their synergism against dermatophytes**
(2010) *Mycoses*, 53 (2), pp. 123-129. [Cited 27 times](#).
doi: 10.1111/j.1439-0507.2008.01676.x
[View at Publisher](#)

☐ Silva, G.H., Teles, H.L., Zanardi, L.M., Marx Young, M.C., Eberlin, M.N., Hadad, R., Pfenning, L.H., (...), Araújo, A.R.
3

Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert](#) | [Set citation feed](#)

Related documents

Bioassay guided isolation of an antidermatophytic active constituent from the stem bark of Entada spiralis Ridl | Pengasingan komponen aktif antiderma secara berperingkat dari kulit pokok Entada spiralis Ridl
Harun, A. , Mat So'Ad, S.Z. , Hassan, N.M.
(2015) *Malaysian Journal of Analytical Sciences*

Studies on chemical constituents in seeds of Entada phaseoloide
Zhang, Y. , Zhang, H.-W. , Zou, Z.-M.
(2008) *Chinese Pharmaceutical Journal*

Triterpene saponins from Entada phaseoloides
Xiong, H. , Mei, Z. , Yang, G.
(2013) *Helvetica Chimica Acta*

[View all related documents based on references](#)

Find more related documents in Scopus based on:

[Authors](#) | [Keywords](#)